

Package ‘ordinalpattern’

August 9, 2023

Type Package

Title Tests Based on Ordinal Patterns

Version 0.2.3

Date 2023-8-6

Author Alexander Duerre [aut, cre],
Alexander Schnurr [aut]

Depends gtools,mvtnorm

Maintainer Alexander Duerre <alexander.duerre@udo.edu>

Description Ordinal patterns describe the dynamics of a time series by looking at the ranks of subsequent observations. By comparing ordinal patterns of two times series, Schnurr (2014) <[doi:10.1007/s00362-013-0536-8](https://doi.org/10.1007/s00362-013-0536-8)> defines a robust and non-parametric dependence measure: the ordinal pattern coefficient. Functions to calculate this and a method to detect a change in the pattern coefficient proposed in Schnurr and Dehling (2017) <[doi:10.1080/01621459.2016.1164706](https://doi.org/10.1080/01621459.2016.1164706)> are provided.

License GPL-2 | GPL-3

ByteCompile true

Encoding UTF-8

NeedsCompilation yes

Repository CRAN

Date/Publication 2023-08-09 10:50:11 UTC

R topics documented:

patternchange	2
patterndependence	4

Index	6
--------------	----------

patternchange *Changepoint Detection Using Ordinal Patterns*

Description

Test for a change in the dependence structure of two time series using ordinal patterns

Usage

```
patternchange(tsx, tsy, h=2, conf.level, weight=TRUE, weightfun=NULL, bn=log(length(tsx)),
             kernel=function(x){return(max(0, 1-abs(x)))})
```

```
## S3 method for class 'change'
plot(x, ...)
```

Arguments

tsx	numeric vector of first univariate time series.
tsy	numeric vector of second univariate time series.
h	numeric value determining the length of ordinal pattern.
conf.level	numerical value indicating the confidence level of the test.
weight	logical value indicating whether one uses weights of the L1 norm or the empirical probability of identical patterns; see details.
weightfun	function which defines the weights given the L1 norm between the patterns if weight=TRUE. If no weight-function is given, the canonical weight function is used; see details.
bn	numerical value determining the bandwidth of the kernel estimator used to estimate the long run variance.
kernel	kernel function for estimating the long run variance.
x	object of class "change"
...	further arguments passed to the internal plotting function (plot).

Details

Given two timeseries `tsx` and `tsy` a cusum type statistic tests whether there is a change in the pattern dependence or not. The test is based on a comparison of patterns of length $h+1$ in `tsx` and `tsy`. One can either choose the number of identical patterns (`weight=FALSE`) or a metric that is defined by the `weightfun` argument to measure the difference between patterns (`weight=TRUE`). If no (`weightfun`) is given, the canonical weightfunction is used, which equals 1 if patterns are identical and 0 if the L1 norm of their difference attains the maximal possible value. The value is linear interpolated in between.

The procedure depends on an estimate of the long run variance. Here a kernel estimator is used. A kernel function and a bandwidth can be set using the arguments `kernel` and `bn`. If none of them is given, the bartlett kernel with a bandwidth of $\log(n)$, where n equals the length of the timeseries, is used.

Value

Object with classes "change" and "htest" containing the following values:

statistic	the value of the test statistic. Under the null the test statistic follows asymptotically a Kolmogorov Smirnov distribution.
p.value	the p-value of the test.
estimate	the estimated time of change.
null.value	the jump height of the at most one change point model, which is under the null hypothesis always 0.
alternative	a character string describing the alternative hypothesis.
method	a characters string describing the test.
trajectory	the cumulative sum on which the tests are based on. Could be used for additional plots.

Author(s)

Alexander Dürre

References

Schnurr, A. (2014): An ordinal pattern approach to detect and to model leverage effects and dependence structures between financial time series, *Statistical Papers*, vol. 55, 919–931.

Schnurr, A., Dehling, H. (2017): Testing for Structural Breaks via Ordinal Pattern Dependence, *Journal of the American Statistical Association*, vol. 112, 706–720.

See Also

Estimation of the pattern dependence is provided by [patterndependence](#).

Examples

```
set.seed(1066)
a1 <- cbind(rnorm(100), rnorm(100))
a2 <- rmvnorm(100, sigma=matrix(c(1, 0.8, 0.8, 1), ncol=2))
A <- rbind(a1, a2)
testresult <- patternchange(A[,1], A[,2])
plot(testresult)
testresult
```

patterndependence *Ordinal Pattern Dependence*

Description

Calculates the standard ordinal pattern coefficient and related values

Usage

```
patterndependence(tsx, tsy, h=2, block=FALSE, first=TRUE)
```

```
## S3 method for class 'pattern'
plot(x, ...)
## S3 method for class 'pattern'
print(x, ...)
```

Arguments

tsx	numeric vector representing the first univariate time series.
tsy	numeric vector representing the second univariate time series.
h	numeric value determining the length of the ordinal pattern.
block	logical value determining whether patterns are calculated on disjoint blocks or overlapping blocks.
first	logical value indicating which observations are dropped if block == TRUE and the time series length is no multiple of h+1.
x	object of class "pattern", which is the output of patterndependence
...	further arguments passed to the internal plotting function.

Details

The standard ordinal pattern coefficient is a non-parametric and robust measure of dependence between two time series. It is based on ordinal patterns, which are defined as sequences of ranks of h+1 subsequent observations. This sequences of subsequent observations can either move one observation per time or a whole block of h+1 observations. The former is preferred since it uses more information. If one chooses the later, one has to decide whether the first or the last observations are removed in case that the time series length is no multiple of h+1.

Beside the standard ordinal pattern coefficient, which range from -1 to 1, one can also look at the positive and negative ordinal pattern coefficient, which roughly measures wheather there are unusual many identical or opposite patterns in the time series.

The plot function draws both time series and shows the six most frequent coinciding pattern with counts on the right. At the bottom, the location of these coinciding patterns is visualized.

Value

Object of class "pattern" containing the following values:

patterncoef	standard ordinal pattern coefficient.
alpha	positive ordinal pattern coefficient, see details.
beta	negative ordinal pattern coefficient, see details.
numbequal	number of equal ordinal patterns.
numbopposite	number of opposite ordinal patterns.
PatternXz	number of ordinal patterns in first time series.
PatternYz	number or ordinal patterns in second time series.
coding	coding of the ordinal patterns, used in PatternXz and PatternYz.
PatternX	numeric vector representing the time series of patterns in tsx.
PatternY	numeric vector representing the time series of patterns in tsy.
tsx	numeric vector representing the first univariate time series.
tsy	numeric vector representing the second univariate time series.
maxpat	number representing the maximal pattern code.
block	logical value determining whether patterns are calculated on disjoint blocks or overlapping blocks.
h	number of consecutive observations defining one pattern
tablesame	numeric vector representing the number of coinciding patterns, apportioned into different patterns.
tablesame	numeric vector representing the number of reflected patterns, apportioned into different patterns.
indexsame	logic vector indicating whether patterns in both time series coincide.
indexsame	logic vector indicating whether patterns in both time series are reflected.

Author(s)

Alexander Dürre

References

Schnurr, A. (2014): An ordinal pattern approach to detect and to model leverage effects and dependence structures between financial time series, *Statistical Papers*, vol. 55, 919–931.

Schnurr, A., Dehling, H. (2017): Testing for Structural Breaks via Ordinal Pattern Dependence, *Journal of the American Statistical Association*, vol. 112, 706–720.

Examples

```
set.seed(1066)
patternobj <- patterndependence(rnorm(100), rnorm(100))
plot(patternobj)
```

Index

patternchange, [2](#)
patterndependence, [3](#), [4](#)
plot.change (patternchange), [2](#)
plot.pattern (patterndependence), [4](#)
print.pattern (patterndependence), [4](#)